Minimum Immunization Requirements for Those Entering a Child Care Facility or School in Illinois, Fall-2016 (Page 1)

Vaccine	Child Care Facility,	Kindergarten through 12 th Grade		Minimum Intervals
Requirement	Preschool, Early Childhood Pre-Kindergarten Programs	First Entry into School (Kindergarten or First Grade)	Other Grades	Allowed Between Doses and Other Options for Proof of Immunity
DTP/DTaP/ or Tdap, Td (Diphtheria, Tetanus, Pertussis)	Three doses by 1 year of age One additional booster dose	Four or more doses of DTP/DTaP with the last dose qualifying as a booster and received on or after the 4 th	Three or more doses of DTP/DTaP or Td; with the last dose qualifying as a booster if received on or after the 4 th birthday For Students entering 6 th thru 12 th	Minimum interval between series doses: 4 weeks (28 days) Between series and booster: 6 months No proof of immunity allowed
	by 2 nd birthday	birthday	grades: 1 dose of Tdap	
Polio	Two doses by 1 year of age. One additional dose by 2 nd birthday	Four or more doses of the same type of Polio vaccine with the last dose qualifying as a booster and received on or after the 4 th birthday. (progressive requirement)	Three or more doses of Polio with the last dose qualifying as a booster and received on or after the 4 th birthday. *If the series is given in any combination of polio vaccine types, 4 or more doses are required with the last being a booster on or after the 4 th birthday.	Minimum interval between series doses: 4 weeks (28 days) For Grade K: 6 month interval between 3 dose series and booster; booster must be on or after 4 th birthday No proof of immunity allowed
Measles	One dose on or after the 1 st birthday but prior to 24 months of age	Two doses of Measles Vaccine , the 1 st dose must have been received on after the 1 st birthday and the second dose no less than 4 weeks (28 days) later.		Laboratory evidence of measles immunity <u>OR</u> Certified physician verification* of measles disease by date of illness *Cases diagnosed after 7/1/2002 must include lab evidence of infection
Rubella	One dose on or after the 1 st birthday but prior to 24 months of age	Two doses of Rubella Vaccine , the 1 st dose must have been received on after the 1 st birthday and the second dose no less than 4 weeks (28 days) later.		Laboratory evidence of rubella immunity History of disease is not acceptable proof of immunity to rubella
Mumps	One dose on or after the 1 st birthday but prior to 24 months of age	Two doses of Mumps Vaccine , the 1 st dose must have been received on after the 1 st birthday and the second dose no less than 4 weeks (28 days) later.		Laboratory evidence of mumps immunity OR Certified physician verification of mumps disease by date of illness
Haemophilus influenzae type b	Refer to Hib ACIP recs for schedule for series Children 24-59 mos. without series must have one dose	Not required after the 5 th birthday (60 months of age)		Refer to ACIP Hib vaccination schedule No proof of immunity allowed
Invasive Pneumococcal Disease	Refer to PCV ACIP recs for schedule for series Children 24-59 mos. without series must have one dose		5 th birthday (60 months of age)	Refer to ACIP PCV vaccination schedule
	after 24 months of age			No proof of immunity allowed

Minimum Immunization Requirements for Those Entering a Child Care Facility or School in Illinois, Fall-2016 (Page 2)

Vaccine	Child Care Facility,	Kindergarten through 12 th Grade		Minimum Intervals
Requirement	Preschool, Early Childhood Pre-Kindergarten Programs	First Entry into School (Kindergarten or First Grade)	All Other Grades	Allowed Between Doses and Other Options for Proof of Immunity
Hepatitis B	Three doses for all children Third dose must have been administered on or after 6 months of age (168 days)	No Requirements	For Students entering grades 6 thru 12: Three doses hepatitis B vaccine administered at recommended intervals	Minimum intervals between doses: 1 & 2- at least 4 weeks (28 days) 2 & 3 - at least 2 months (56 days) 1 & 3 - at least 4 months (112 days) Laboratory evidence of prior or current infection
Varicella (progressive requirement)	One dose on or after 1 st birthday	Two doses of Varicella Vaccine, for Students entering Kindergarten or first grade The 1 st dose must have been received on after the 1 st birthday and the second dose no less than 4 weeks (28 days) later.	One dose of Varicella on or after the 1 st birthday for Students entering grades 3, 4, 5, & 12	Minimum intervals for administration: The 1 st dose must have been received on after the 1 st birthday and the second dose no less than 4 weeks (28 days) later. Statement from physician or health care provider verifying disease history <u>OR</u> Laboratory evidence of varicella immunity
			Two doses of Varicella Vaccine for Students entering grades 2, 6, 7, 8, 9, 10, & 11	
Meningococcal Conjugate Vaccine (progressive requirement)	No Requirements	No Requirements	Applies to Students entering grades 6, 7, or 12 beginning 2016-2017 school year 1 dose of Meningococcal vaccine for entry to grade 6 or 7 2 doses of Meningococcal vaccine at	Minimum intervals for administration: The first dose received on or after the 10 th birthday; second dose at least eight weeks after the 1 st dose. Only one dose is required if the first dose was received at 16 years
			entry to 12 TH grade	of age or older. No proof of immunity allowed.

Source: Child Health Examination Code/Part 665

Prepared by: Illinois Department of Public Health Immunization Section, January 2016