

Recommended Vaccines

	Child Care Centers (by 12 Months of Age)	Kindergarten through 5th Grade	6th through 12th Grade
DTaP	4*	5*	5*
IPV	3*	4*	4*
MMR	1*	2*	2*
Varicella	1*	1*	2**
Hib	4*		
PCV	4*		
Tdap			1*
Rota	3		
Hep A	1	2	2
HepB	3	3	3
HPV			2
MCV			2

Prior to entering any Alabama school or child care center, an up-to-date Certificate of Immunization must be submitted for each child.

*Code of Alabama §16-30-4 **1 dose for <13 years of age or 2 doses for >13 years of age



For more information,
 visit alabamapublichealth.gov/immunization
 or facebook.com/AlabamaImmunizationInfo



Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	19-23 months	2-3 years	4-6 years
HepB	HepB			HepB						
	RV	RV	RV							
	DTap	DTap	DTap		DTap					DTap
	Hib	Hib	Hib	Hib						
	PCV13	PCV13	PCV13	PCV13						
	IPV	IPV	IPV							
				Influenza (yearly)*						
					MMR					MMR
					Varicella					Varicella
	HepA [§]									

■ Range of recommended ages for all children

*Two doses given at least four weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.

§Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 months after the last dose. HepA vaccination may be given to any child 12 months and older to protect against hepatitis A. Children and adolescents who did not receive the HepA vaccine and are at high risk should be vaccinated against hepatitis A.

	7-8 years	9-10 years	11-12 years	13-15 years	16-18 years
Flu Influenza ¹	■				
Tdap (Tetanus, diphtheria, pertussis) ²	■		■	■	
HPV (Human papillomavirus) ³		■	■	■	
Meningococcal MenACWY ⁴	■		■	■	■
Meningococcal MenB ⁵		■			■
Pneumococcal	■				
Hepatitis B	■				
Hepatitis A	■				
Polio	■				
MMR (Measles, mumps, rubella)	■				
Chickenpox Varicella	■				

1. Everyone 6 months and older should get a flu vaccine every year.
2. All 11-12 year olds should get one shot of Tdap.
3. All 11-12 year olds should get a 2-shot series of HPV vaccine. A 3-shot series is needed for those with weakened immune systems and those who start the series at 15 years or older.
4. All 11-12 year olds should get one shot of meningococcal conjugate (MenACWY). A booster shot is recommended at age 16.
5. Teens 16-18 years old may be vaccinated with a serogroup B meningococcal (MenB) vaccine.

These shaded boxes indicate when the vaccine is recommended for all children unless your doctor tells you that your child cannot safely receive the vaccine.

These shaded boxes indicate the vaccine should be given if a child is catching up on missed vaccines.

These shaded boxes indicate when the vaccine is recommended for children with certain health or lifestyle conditions that put them at risk for serious disease. See vaccine-specific recommendations at [cdc.gov/vaccines/hcp/acip-recs/index.html](https://www.cdc.gov/vaccines/hcp/acip-recs/index.html).

This shaded box indicates children not at increased risk may get the vaccine if they wish after speaking to a provider.

[cdc.gov/vaccines/schedules/easy-to-read/child-easyread.html#table-child](https://www.cdc.gov/vaccines/schedules/easy-to-read/child-easyread.html#table-child)

[cdc.gov/vaccines/schedules/easy-to-read/adolescent-easyread.html](https://www.cdc.gov/vaccines/schedules/easy-to-read/adolescent-easyread.html)