

	TERM 1		
Term 1 Dates	MS College and Career Readiness Standards		
WEEK 1-2	Number Sense (NS)		
August 6-16, 2024	<ul> <li>7.NS.1 Apply and extend previous understandings of addition and subtraction to add and subtract rational numbers; represent addition and subtraction on a horizontal or vertical number line diagram.</li> <li>a. Describe situations in which opposite quantities combine to make 0. For example, a hydrogen atom has 0 charge because its two constituents are oppositely charged.</li> <li>b. Understand p + q as the number located a distance  q  from p, in the positive or negative direction depending on whether q is positive or negative. Show that a number and its opposite have a sum of 0 (are additive inverses). Interpret sums of rational numbers by describing real world contexts.</li> <li>c. Understand subtraction of rational numbers as adding the additive inverse, p - q = p + (-q). Show that the distance between two rational numbers on the number line is the absolute value of their difference, and apply this principle in real-world contexts.</li> <li>d. Apply properties of operations as strategies to add and subtract rational numbers.</li> </ul>		
WEEK 3	<b>7.NS.2</b> Apply and extend previous understandings of multiplication and division and of fractions to multiply		
August 19 -23, 2024	and divide rational numbers.  a. Understand that multiplication is extended from fractions to rational numbers by requiring that operations continue to satisfy the properties of operations, particularly the distributive property, leading to products such as (-1)(-1) = 1 and the rules for multiplying signed numbers. Interpret products of rational numbers by describing real-world contexts.  b. Understand that integers can be divided, provided that the divisor is not zero, and every quotient of integers (with non-zero divisor) is a rational number. If p and q are integers, then - (p/q) = (-p)/q = p/(-q). Interpret quotients of rational numbers by describing real world contexts.  c. Apply properties of operations as strategies to multiply and divide rational numbers.  d. Convert a rational number to a decimal using long division; know that the decimal form of a rational number terminates in 0s or eventually repeats.		
WEEK 4	<b>7.NS.3</b> Solve real-world and mathematical problems involving the four operations with rational numbers.		
August 26-30, 2024			
WEEK 5	Expressions and Equations (EE)		



September 3-6, 2024	<b>7.EE.1</b> Apply properties of operations as strategies to add, subtract, factor, and expand linear expressions with		
	rational coefficients.		
WEEK 6	<b>7.EE.2</b> Understand that rewriting an expression in different forms in a problem context can shed light on the		
September 9-13,	problem and how the quantities in it are related. For example, a + 0.05a = 1.05a means that "increase by 5%"		
2024	is the same as "multiply by 1.05."		
WEEK 7-8	7.EE.3 Solve multi-step real-life and mathematical problems posed with positive and negative rational		
September 16-27,	numbers in any form (whole numbers, fractions, and decimals), using tools strategically. Apply properties of		
2024	operations to calculate with numbers in any form; convert between forms as appropriate; and assess the		
	reasonableness of answers using mental computation and estimation strategies. For example: If a woman		
	making \$25 an hour gets a 10% raise, she will make an additional 1/10 of her salary an hour, or \$2.50, for a		
	new salary of \$27.50. If you want to place a towel bar 9 3/4 inches long in the center of a door that is 27 1/2		
	inches wide, you will need to place the bar about 9 inches from each edge; this estimate can be used as a		
	check on the exact computation. 3		
WEEK 9	Review Term 1 Standards		
September 30 -			
October 4, 2024			
WEEK 10	BMAs / First 9 Weeks Assessment (End of Term 1)		
October 7-11, 2024			

IERIVI	Т	Е	R	M	1
--------	---	---	---	---	---

### **Recurring Standards**

(Standards that have been introduced but will be reviewed or revisited during the nine weeks.)

#### Standards:

**6.NS.3** Add, subtract, multiply, and divide multi-digit decimals using the standard algorithm for each operation.

**6.NS.5** Understand that positive and negative numbers are used together to describe quantities having opposite directions or values.

**6.NS.7** Understand ordering and absolute value of rational numbers.



,		TERM 2		
equations and inequalities to solve problems by reasoning about the quantities.  a. Solve word problems leading to equations of the form px + q = r and p(x + q) = r, where p, q, and r are specific rational numbers. Solve equations of these forms fluently. Compare an algebraic solution to an arithmetic solution, identifying the sequence of the operations used in each approach. For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?  b. Solve word problems leading to inequalities of the form px + q > r or px + q < r, where p, q, and r are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.  WEEK 3  October 28 -  November 1, 2024  Ratio and Proportion (RP)  7.RP.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour  WEEK 4  November 4-15, 2024  7.RP. 2 Recognize and represent proportional relationships between quantities. a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin. b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships by equations. For example, if total cost t is proportional to the number n of items can be expressed as t = pn. d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and	Term 2 Dates	MS College and Career Readiness Standards		
a. Solve word problems leading to equations of the form px + q = r and p(x + q) = r, where p, q, and r are specific rational numbers. Solve equations of these forms fluently. Compare an algebraic solution to an arithmetic solution, identifying the sequence of the operations used in each approach. For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?  b. Solve word problems leading to inequalities of the form px + q > r or px + q < r, where p, q, and r are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.  WEEK 3  October 28 - November 1, 2024  November 1, 2024  TARP.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour  WEEK 4  November 4-15, 2024  7.RP.2 Recognize and represent proportional relationships between quantities.  a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.  b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.  c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple	WEEK 1-2	7.EE.4 Use variables to represent quantities in a real-world or mathematical problem, and construct simple		
specific rational numbers. Solve equations of these forms fluently. Compare an algebraic solution to an arithmetic solution, identifying the sequence of the operations used in each approach. For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?  b. Solve word problems leading to inequalities of the form px + q > r or px + q < r, where p, q, and r are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.  WEEK 3  October 28 - November 1, 2024  7.RP.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour  WEEK 4  November 4-15, 2024  7.RP. 2 Recognize and represent proportional relationships between quantities. a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin. b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships. c. Represent proportional relationships be quations. For example, if total cost t is proportional to the number of items can be expressed as t = pn. d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple	October 15-25, 2024	equations and inequalities to solve problems by reasoning about the quantities.		
arithmetic solution, identifying the sequence of the operations used in each approach. For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?  b. Solve word problems leading to inequalities of the form px + q > r or px + q < r, where p, q, and r are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.  Ratio and Proportion (RP)  7.RP.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour  7.RP.2 Recognize and represent proportional relationships between quantities.  a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.  b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships by equations. For example, if total cost t is proportional to the number n of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple				
perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?  b. Solve word problems leading to inequalities of the form px + q > r or px + q < r, where p, q, and r are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.  WEEK 3  October 28 - November 1, 2024  October 28 - November 1, 2024  WEEK 4  November 4-15, 2024  7.RP. 2 Recognize and represent proportional relationships between quantities.  A Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.  b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.  c. Represent proportional relationships.  c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple		, , , , , , , , , , , , , , , , , , , ,		
specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.  Ratio and Proportion (RP)  7.RP.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour  WEEK 4  November 4-15, 2024  7.RP.2 Recognize and represent proportional relationships between quantities. a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin. b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships by equations. For example, if total cost t is proportional to the number of items can be expressed at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn. d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple				
problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.  Ratio and Proportion (RP) 7.RP.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour  WEEK 4 November 4-15, 2024  7.RP. 2 Recognize and represent proportional relationships between quantities. a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin. b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships. c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn. d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple		b. Solve word problems leading to inequalities of the form $px + q > r$ or $px + q < r$ , where p, q, and r are		
your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.  Ratio and Proportion (RP)  7.RP.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour  7.RP. 2 Recognize and represent proportional relationships between quantities.  a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.  b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.  c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple		specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the		
the solutions.  Ratio and Proportion (RP)  7.RP.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour  WEEK 4  November 4-15, 2024  7.RP. 2 Recognize and represent proportional relationships between quantities. a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin. b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships. c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn. d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple		problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want		
WEEK 3 October 28 - November 1, 2024 November 1, 2024 November 1, 2024  WEEK 4 November 4-15, 2024 November 4-15, 2024 November 5  Ratio and Proportion (RP)  7.RP. 2 Recognize and represent proportional relationships between quantities. A Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through descriptions of proportional relationships.  C. Represent proportional relationships. C. Represent proportional relationships between the total cost t is proportional to the number n of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP. 3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple				
<ul> <li>7.RP.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour</li> <li>7.RP. 2 Recognize and represent proportional relationships between quantities.         <ul> <li>a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.</li> <li>b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.</li> <li>c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn.</li> <li>d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of</li> </ul> </li> <li>WEEK 5</li> <li>7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple</li> </ul>				
November 1, 2024  quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour  7.RP. 2 Recognize and represent proportional relationships between quantities.  a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.  b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.  c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple				
<ul> <li>compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour</li> <li>WEEK 4</li> <li>November 4-15,</li> <li>2024</li> <li>a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.         <ul> <li>b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.</li> <li>c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn.</li> <li>d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of</li> </ul> </li> <li>WEEK 5</li> <li>7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple</li> </ul>				
<ul> <li>7.RP. 2 Recognize and represent proportional relationships between quantities.         <ul> <li>a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.</li></ul></li></ul>	November 1, 2024			
a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.  b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.  c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple				
in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.  b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.  c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple				
the origin.  b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.  c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple				
descriptions of proportional relationships.  c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple	2024			
c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn. d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple				
number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn.  d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple				
number of items can be expressed as t = pn. d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple				
d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of  WEEK 5  7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple		· · · · · ·		
WEEK 5 7.RP.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple		· · ·		
	WEEK E			
inoveninci 10-22, Initerest, tax, markups and markuowns, gratuities and commissions, fees, percent increase and decrease,				
	2024	, , , , , , , , , , , , , , , , , , , ,		



WEEK 6	GEOMETRY (G)		
Dec. 2-6, 2024	<b>7.G.1</b> Solve problems involving scale drawings of geometric figures, including computing actual lengths and		
	areas from a scale drawing and reproducing a scale drawing at a different scale.		
WEEK 7	7.G.2 Draw (freehand, with ruler and protractor, and with technology) geometric shapes with given		
Dec. 9-13, 2024	conditions. Focus on constructing triangles from three measures of angles or sides, noticing when the		
	conditions determine a unique triangle, more than one triangle, or no triangle.		
	7.G.3 Describe the two-dimensional figures that result from slicing three-dimensional figures, as in plane		
	sections of right rectangular prisms and right rectangular pyramids.		
WEEK 8	Review Term 2 Standards		
Dec. 9-13, 2024			
WEEK 9	BMAs / Second 9 Weeks Assessment (End of Term 2)		
December 16-20,			
2024			

# **TERM 2**

### **Recurring Standards**

(Standards that have been introduced but will be reviewed or revisited during the nine weeks.)

#### Standards:

7.NS.1, 7.NS.2, 7.NS.3, and 7.EE.3



WEEK 2 January 13-17, 2025 WEEK 2 January 13-17, 2025 WEEK 3 January 13-17, 2025  WEEK 3 January 13-17, 2025  WEEK 3 January 13-17, 2025  WEEK 3 January 21-31, 2025  WEEK 3 JANUARY 3-7, 2025  WEEK 4 JANUARY 3-7, 2025  WEEK 4 JANUARY 3-7, 2025  WEEK 5 JANUARY 3-7, 2025  WEEK 6  WEEK 6		TERM 3			
WEEK 2 January 13-17, 2025 write and solve simple equations for an unknown angle in a figure.  7.6.4 Know the formulas for the area and circumference of a circle and use them to solve problems; give an informal derivation of the relationship between the circumference and area of a circle.  7.6.6 Solve real-world and mathematical problems involving area, volume and surface area of two- and three-dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms.  STATISTICS and PROBABILITY (SP)  7.5.9.1 Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences.  7.5.9.2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  7.5.9.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.5.9.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a	Term 3 Dates	MS College and Career Readiness Standards			
### A comparison of the formulas for the area and circumference of a circle and use them to solve problems; give an informal derivation of the relationship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of two and three-dimensionship between the circumference and area of a circle.  ### A comparison of the relationship between the circumference and area of a circle.  ### A comparison of two and three-dimensionship between the circumference and area of a circle.  ### A comparison of the prison of the polymers on the sample of two population by examining a sample of the polymers on the base of the polymers on the analysis of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  #### A comparison of the polymers of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of player	WEEK 1				
Informal derivation of the relationship between the circumference and area of a circle.  WEEK 3-4 January 21 - 31, 2025  WEEK 5 February 3-7, 2025  TAISTICS and PROBABILITY (SP) 7.SP-1 Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences.  7.SP-2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  WEEK 6 February 10-14, 2025  WEEK 6 February 10-14, 2025  T.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.	· · · · · · · · · · · · · · · · · · ·				
7.G.6 Solve real-world and mathematical problems involving area, volume and surface area of two- and three-dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms.  STATISTICS and PROBABILITY (SP)  7.SP.1 Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences.  7.SP.2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  WEEK 6  February 10-14, 2025  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.		• =			
January 21 - 31, 2025 dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms.  STATISTICS and PROBABILITY (SP)  7.SP.1 Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences.  7.SP.2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  WEEK 6  February 10-14, 2025  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.	· · · · · · · · · · · · · · · · · · ·	·			
February 3-7, 2025  Februa	_	i i i i i i i i i i i i i i i i i i i			
7.SP.1 Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences.  7.SP.2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.	January 21 - 31, 2025	dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms.			
the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences.  7.SP.2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.	WEEK 5	STATISTICS and PROBABILITY (SP)			
of that population. Understand that random sampling tends to produce representative samples and support valid inferences.  7.SP.2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  WEEK 6  February 10-14, 2025  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.	February 3-7, 2025	<b>7.SP.1</b> Understand that statistics can be used to gain information about a population by examining a sample of			
valid inferences.  7.SP.2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  WEEK 6  February 10-14, 2025  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.		the population; generalizations about a population from a sample are valid only if the sample is representative			
7.SP.2 Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.		of that population. Understand that random sampling tends to produce representative samples and support			
of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  WEEK 6 February 10-14, 2025 February 10-14, 2025  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.		valid inferences.			
estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  WEEK 6 February 10-14, 2025  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.		<b>7.SP.2</b> Use data from a random sample to draw inferences about a population with an unknown characteristic			
from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  WEEK 6  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.		of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in			
from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.  WEEK 6  7.SP.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.		estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words			
WEEK 6 February 10-14, 2025 Fe		from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far			
variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.					
variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.	WEEK 6	i s			
variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.	February 10-14, 2025				
height of players on the soccer team, about twice the variability on either team; on a dot plot, the separation between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.	, ,				
between the two distributions of heights is noticeable.  7.SP.4 Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.		, , , , , , , , , , , , , , , , , , , ,			
<b>7.SP.4</b> Use measures of center and measures of variability (i.e. inter-quartile range) for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.					
random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.					
whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.		. ,			
chapter of a fourth-grade science book.					
·		, , , , , , , , , , , , , , , , , , , ,			
WEEK / LASE 5 Understand that the probability of a chance event is a number between 0 and 1 that expresses the	WEEK 7	<b>7.SP.5</b> Understand that the probability of a chance event is a number between 0 and 1 that expresses the			
	February 18-21, 2025				



	unlikely event, a probability around 1/2 indicates an event that is neither unlikely nor likely, and a probability near 1 indicates a likely event.	
WEEK 8	Review Term 3 Standards	
February 24-28, 2025		
WEEK 9	BMAs / Third 9 Weeks Assessment (End of Term 3)	
March 3-7, 2025		

	TERM 3			
	Recurring Standards			
	(Standards that have been introduced but will be reviewed or revisited during the nine weeks.)			
Standards:				
4.MD.7	6.G.1	6.G.4	6.SP.1	
4.G.1	6.G.2	7.G.2	6.SP.5	



	TERM 4
Term 4 Dates	MS College and Career Readiness Standards
WEEK 1	<b>7.SP.6</b> Approximate the probability of a chance event by collecting data on the chance process that produces it
March 17-21, 2025	and observing its long-run relative frequency, and predict the approximate relative frequency given the probability. For example, when rolling a number cube 600 times, predict that a 3 or 6 would be rolled roughly 200 times, but probably not exactly 200 times.
	7.SP.7 Develop a probability model and use it to find probabilities of events. Compare probabilities from a model to observed frequencies; if the agreement is not good, explain possible sources of the discrepancy. a. Develop a uniform probability model by assigning equal probability to all outcomes, and use the model to determine probabilities of events. For example, if a student is selected at random from a class, find the probability that Jane will be selected and the probability that a girl will be selected. b. Develop a probability model (which may not be uniform) by observing frequencies in data generated from a chance process. For example, find the approximate probability that a spinning penny will land heads up or that a tossed paper cup will land open-end down. Do the outcomes for the spinning penny appear to be equally likely based on the observed frequencies?
WEEK 2	<b>7.SP.8</b> Find probabilities of compound events using organized lists, tables, tree diagrams, and simulation.
March 24-28, 2025	a. Understand that, just as with simple events, the probability of a compound event is the fraction of outcomes in the sample space for which the compound event occurs.  b. Represent sample spaces for compound events using methods such as organized lists, tables and tree diagrams. For an event described in everyday language (e.g., "rolling double sixes"), identify the outcomes in the sample space which compose the event.
	c. Design and use a simulation to generate frequencies for compound events. For example, use random digits as a simulation tool to approximate the answer to the question: If 40% of donors have type A blood, what is the probability that it will take at least 4 donors to find one with type A blood?
WEEK 3	Review Statistics and Probability (SP) Standards
March 31-April 4, 2025	
<b>WEEK 4</b> April 7-11, 2025	Review Number Sense (NS), Ratio and Proportion (RP) Standards
<b>WEEK 5</b> April 14-17, 2025	Review Expressions and Equations (EE), Geometry (G) Standards



WEEK 6	Review all standards (Practice Test)		
April 22-25, 2025			
WEEK 7-8	State Testing		
WEEK 9	BMA's / Fourth 9 weeks Assessments (End of Term 4)		

# TERM 4

## **Recurring Standards**

(Standards that have been introduced but will be reviewed or revisited during the nine weeks.)

Standards:

7.SP.1, 7.SP.2, and 7.SP.5